arXiv:math/0411635v1 [math.QA] 29 Nov 2004

On algebras of gauge transformations in a general setting

G.Sardanashvily
Department of Theoretical Physics, Moscow State University, 117234 Moscow, Russia

Abstract We consider a Lagrangian system on a fiber bundle and its gauge transformations depending on derivatives of dynamic variables and gauge parameters of arbitrary order. We say that gauge transformations form an algebra if they generate a nilpotent BRST operator.

1 Introduction

In a general setting, one can say that a family of gauge transformations depending on parameters is an algebra if the Lie bracket of arbitrary two gauge transformations depending on different parameter functions is again a gauge transformation depending on some parameter function. The goal is to formulate this condition in strict mathematical terms. For instance, gauge transformations in gauge theory on a principal bundle form a finite-dimensional real (or complex) Lie algebra. Gauge transformations of a certain class of field theories constitute sh-Lie algebras [4]. It may happen that gauge transformations are not assembled into an algebra or form an algebra on-shell [6].

We consider a Lagrangian system on a smooth fiber bundle $Y \to X$ subject to gauge transformations depending both on derivatives of dynamic variables of arbitrary order and a finite family of gauge parameters and their derivatives of arbitrary order.

Let J^rY, $r = 1, \ldots$, be finite order jet manifolds of sections of $Y \to X$. In the sequel, the index $r = 0$ stands for Y. Given bundle coordinates (x^λ, y^i) on Y, jet manifolds J^rY are endowed with the adapted coordinates $(x^\lambda, y^i, y^i_\Lambda)$, where $\Lambda = (\lambda_k...\lambda_1)$, $k = 1, \ldots, r$, is a symmetric multi-index. We use the notation $\lambda + \Lambda = (\lambda\lambda_k...\lambda_1)$ and

$$d_\lambda = \partial_\lambda + \sum_{0\leq|\Lambda|} y^i_{\lambda+\Lambda}\partial^\Lambda_i, \qquad d_\Lambda = d_{\lambda_r}\circ\cdots\circ d_{\lambda_1}. \tag{1}$$

In order to describe gauge transformations depending on parameters, let us consider Lagrangian formalism on the bundle product

$$E = Y \underset{X}{\times} V, \tag{2}$$

where $V \to X$ is a vector bundle whose sections are gauge parameter functions [2]. Let $V \to X$ be coordinated by (x^λ, ξ^r). Then gauge transformations are represented by a differential operator

$$\upsilon = \sum_{0\leq|\Lambda|\leq m} \upsilon^{i,\Lambda}_r(x^\lambda, y^i_\Sigma)\xi^r_\Lambda\partial_i \tag{3}$$

on E (2) which is linear on V and takes its values into the vertical tangent bundle VY of $Y \to X$. Given a section $\xi(x)$ of $V \to X$, the pull-back

$$\xi^*\upsilon = \sum_{0\leq|\Lambda|\leq m} \upsilon_r^{i,\Lambda}(x^\lambda, y^i_\Sigma) d_\Lambda \xi^r(x)\partial_i \tag{4}$$

of υ (3) onto Y is a gauge transformation depending on a parameter function $\xi(x)$.

By means of a replacement of even gauge parameters ξ^r and their jets ξ^r_Λ with the odd ghosts c^r and their jets c^r_Λ, the operator (3) defines a graded derivation

$$\upsilon = \sum_{0\leq|\Lambda|\leq m} \upsilon_r^{i,\Lambda}(x^\lambda, y^i_\Sigma) c^r_\Lambda \partial_i \tag{5}$$

of the algebra of the original even fields and odd ghosts. Its extension

$$\upsilon = \sum_{0\leq|\Lambda|\leq m} \upsilon_r^{i,\Lambda} c^r_\Lambda \partial_i + u^r\partial_r \tag{6}$$

to ghosts is called the BRST transformation if it is nilpotent.

We say that gauge transformations (4) make up an algebra if they generate a BRST transformation (6). One can think of the nilpotency conditions (36) – (37) as being the generalized commutation relations and Jacobi identity, respectively. This definition is especially convenient for BRST theory and BV quantization [1].

2 Gauge systems on fiber bundles

In Lagrangian formalism on a fiber bundle $Y \to X$, Lagrangians and their Euler–Lagrange operators are represented by elements of the following graded differential algebra (henceforth GDA).

With the inverse system of jet manifolds

$$\begin{tikzcd} X & Y \arrow[l, "\pi"'] & J^1Y \arrow[l, "\pi^1_0"'] & \cdots J^{r-1}Y \arrow[l] & J^rY \arrow[l, "\pi^r_{r-1}"'] & \cdots, \arrow[l] \end{tikzcd} \tag{7}$$

one has the direct system

$$\begin{tikzcd} \mathcal{O}^*X \arrow[r, "\pi^*"] & \mathcal{O}^*Y \arrow[r, "\pi^1_0{}^*"] & \mathcal{O}^*_1Y \arrow[r] & \cdots \mathcal{O}^*_{r-1}Y \arrow[r, "\pi^r_{r-1}{}^*"] & \mathcal{O}^*_rY \arrow[r] & \cdots \end{tikzcd} \tag{8}$$

of GDAs $\mathcal{O}^*_rY$ of exterior forms on jet manifolds J^rY with respect to the pull-back monomorphisms $\pi^r_{r-1}{}^*$. Its direct limit $\mathcal{O}^*_\infty Y$ is a GDA consisting of all exterior forms on finite order jet manifolds modulo the pull-back identification.

The projective limit $(J^\infty Y, \pi^\infty_r : J^\infty Y \to J^rY)$ of the inverse system (7) is a Fréchet manifold. A bundle atlas $\{(U_Y; x^\lambda, y^i)\}$ of $Y \to X$ yields the coordinate atlas

$$\{((\pi^\infty_0)^{-1}(U_Y); x^\lambda, y^i_\Lambda)\}, \qquad {y'}^i_{\lambda+\Lambda} = \frac{\partial x^\mu}{\partial x'^\lambda} d_\mu {y'}^i_\Lambda, \qquad 0 \leq |\Lambda|, \tag{9}$$

of $J^\infty Y$, where d_μ are the total derivatives (1). Then $\mathcal{O}^*_\infty Y$ can be written in a coordinate form where the horizontal one-forms $\{dx^\lambda\}$ and the contact one-forms $\{\theta^i_\Lambda = dy^i_\Lambda - y^i_{\lambda+\Lambda}dx^\lambda\}$ are generating elements of the $\mathcal{O}^0_\infty U_Y$-algebra $\mathcal{O}^*_\infty U_Y$.

There is the canonical decomposition $\mathcal{O}^*_\infty Y = \oplus \mathcal{O}^{k,m}_\infty Y$ of $\mathcal{O}^*_\infty Y$ into $\mathcal{O}^0_\infty Y$-modules $\mathcal{O}^{k,m}_\infty Y$ of k-contact and m-horizontal forms together with the corresponding projectors $h_k : \mathcal{O}^*_\infty Y \to \mathcal{O}^{k,*}_\infty Y$ and $h^m : \mathcal{O}^*_\infty Y \to \mathcal{O}^{*,m}_\infty Y$. Accordingly, the exterior differential on $\mathcal{O}^*_\infty Y$ is split into the sum $d = d_H + d_V$ of the nilpotent total and vertical differentials

$$d_H(\phi) = dx^\lambda \wedge d_\lambda(\phi), \qquad d_V(\phi) = \theta^i_\Lambda \wedge \partial^\Lambda_i \phi, \qquad \phi \in \mathcal{O}^*_\infty Y.$$

Any finite order Lagrangian

$$L = \mathcal{L}\omega : J^r Y \to \overset{n}{\wedge} T^*X, \qquad \omega = dx^1 \wedge \cdots \wedge dx^n, \qquad n = \dim X, \tag{10}$$

is an element of $\mathcal{O}^{0,n}_\infty Y$, while

$$\delta L = \mathcal{E}_i \theta^i \wedge \omega = \sum_{0\leq|\Lambda|} (-1)^{|\Lambda|} d_\Lambda(\partial^\Lambda_i \mathcal{L})\theta^i \wedge \omega \in \mathcal{O}^{1,n}_\infty Y \tag{11}$$

is its Euler–Lagrange operator taking values into the vector bundle

$$T^*Y \underset{Y}{\wedge} (\overset{n}{\wedge} T^*X) = V^*Y \underset{Y}{\otimes} \overset{n}{\wedge} T^*X. \tag{12}$$

A Lagrangian system on a fiber bundle $Y \to X$ is said to be a gauge theory if its Lagrangian L admits a family of variational symmetries parameterized by elements of a vector bundle $V \to X$ and its jet manifolds as follows.

Let $\mathfrak{d}\mathcal{O}^0_\infty Y$ be the $\mathcal{O}^0_\infty Y$-module of derivations of the $\mathbb{R}$-ring $\mathcal{O}^0_\infty Y$. Any $\vartheta \in \mathfrak{d}\mathcal{O}^0_\infty Y$ yields the graded derivation (the interior product) $\vartheta\rfloor\phi$ of the GDA $\mathcal{O}^*_\infty Y$ given by the relations

$$\begin{aligned}
&\vartheta\rfloor df = \vartheta(f), \qquad f \in \mathcal{O}^0_\infty Y,\\
&\vartheta\rfloor(\phi \wedge \sigma) = (\vartheta\rfloor\phi) \wedge \sigma + (-1)^{|\phi|}\phi \wedge (\vartheta\rfloor\sigma), \qquad \phi, \sigma \in \mathcal{O}^*_\infty Y,
\end{aligned}$$

and its derivation (the Lie derivative)

$$\begin{aligned}
&\mathbf{L}_\vartheta \phi = \vartheta\rfloor d\phi + d(\vartheta\rfloor\phi), \qquad \phi \in \mathcal{O}^*_\infty Y, \\
&\mathbf{L}_\vartheta(\phi \wedge \phi') = \mathbf{L}_\vartheta(\phi) \wedge \phi' + \phi \wedge \mathbf{L}_\vartheta(\phi').
\end{aligned} \tag{13}$$

Relative to an atlas (9), a derivation $\vartheta \in \mathfrak{d}\mathcal{O}^0_\infty$ reads

$$\vartheta = \vartheta^\lambda \partial_\lambda + \vartheta^i \partial_i + \sum_{|\Lambda|>0} \vartheta^i_\Lambda \partial^\Lambda_i, \tag{14}$$

where $\{\partial_\lambda, \partial_i^\Lambda\}$ is the dual to the basis $\{dx^\lambda, dy^i_\Lambda\}$ with respect to the interior product $\rfloor$ [5].

A derivation ϑ is called contact if the Lie derivative $\mathbf{L}_\vartheta$ (13) preserves the contact ideal of the GDA $\mathcal{O}^*_\infty Y$ generated by contact forms. A derivation υ (14) is contact iff

$$\vartheta^i_\Lambda = d_\Lambda(\vartheta^i - y^i_\mu\vartheta^\mu) + y^i_{\mu+\Lambda}\vartheta^\mu, \qquad 0 < |\Lambda|. \tag{15}$$

Any contact derivation admits the horizontal splitting

$$\vartheta = \vartheta_H + \vartheta_V = \vartheta^\lambda d_\lambda + (\upsilon^i\partial_i + \sum_{0<|\Lambda|} d_\Lambda \upsilon^i \partial_i^\Lambda), \qquad \upsilon^i = \vartheta^i - y^i_\mu\vartheta^\mu. \tag{16}$$

Its vertical part ϑ_V is completely determined by the first summand

$$\upsilon = \upsilon^i(x^\lambda, y^i_\Lambda)\partial_i, \qquad 0 \leq |\Lambda| \leq k. \tag{17}$$

This is a section of the pull-back $VY \underset{Y}{\times} J^kY \to J^kY$, i.e., a k-order VY-valued differential operator on Y. One calls υ (17) a generalized vector field on Y.

Proposition 1. The Lie derivative of a Lagrangian L (10) along a contact derivation ϑ (16) fulfills the first variational formula

$$\mathbf{L}_\vartheta L = \upsilon\rfloor\delta L + d_H(h_0(\vartheta\rfloor\Xi_L)) + \mathcal{L}d_V(\vartheta_H\rfloor\omega), \tag{18}$$

where Ξ_L is a Lepagean equivalent of L [5]

A contact derivation ϑ (16) is called variational if the Lie derivative (18) is d_H-exact, i.e., $\mathbf{L}_\vartheta L = d_H\sigma$, $\sigma \in \mathcal{O}^{0,n-1}_\infty$. A glance at the expression (18) shows that: (i) ϑ (16) is variational only if it is projected onto X; (ii) ϑ is variational iff its vertical part ϑ_V is well; (iii) it is variational iff $\upsilon\rfloor\delta L$ is d_H-exact.

By virtue of item (ii), we can restrict our consideration to vertical contact derivations $\vartheta = \vartheta_V$. A generalized vector field υ (17) is called a variational symmetry of a Lagrangian L if it generates a variational contact derivation.

Turn now to the notion of a gauge symmetry [2]. Let us consider the bundle product E (2) coordinated by (x^λ, y^i, ξ^r). Given a Lagrangian L on Y, let us consider its pull-back, say again L, onto E. Let ϑ_E be a contact derivation of the $\mathbb{R}$-ring $\mathcal{O}^0_\infty E$, whose restriction

$$\vartheta = \vartheta_E|_{\mathcal{O}^0_\infty Y} = \sum_{0\leq|\Lambda|} d_\Lambda \upsilon^i \partial_i^\Lambda \tag{19}$$

to $\mathcal{O}^0_\infty Y \subset \mathcal{O}^0_\infty E$ is linear in coordinates ξ^r_Ξ. It is determined by a generalized vector field υ_E on E whose projection

$$\upsilon : J^kE \xrightarrow{\upsilon_E} VE \to E \underset{Y}{\times} VY$$

is a linear VY-valued differential operator υ (3) on E. Let ϑ_E be a variational symmetry of a Lagrangian L on E, i.e.,

$$\upsilon_E\rfloor\delta L = \upsilon\rfloor\delta L = d_H\sigma. \qquad (20)$$

Then one says that υ (3) is a gauge symmetry of a Lagrangian L.

Note that any differential operator υ (3) defines a generalized vector field $\upsilon_E = \upsilon$ on E which lives in VY and, consequently, generates a contact derivation $\vartheta_E = \vartheta$ (19).

3 Graded Lagrangian systems

In order to introduce a BRST operator, let us consider Lagrangian systems of even and odd variables. We describe odd variables and their jets on a smooth manifold X as generating elements of the structure ring of a graded manifold whose body is X [5, 8]. This definition reproduces the heuristic notion of jets of ghosts in the field-antifield BRST theory [1, 3].

Recall that any graded manifold $(\mathfrak{A}, X)$ with a body X is isomorphic to the one whose structure sheaf $\mathfrak{A}_Q$ is formed by germs of sections of the exterior product

$$\wedge Q^* = \mathbb{R} \underset{X}{\oplus} Q^* \underset{X}{\oplus} \overset{2}{\wedge} Q^* \underset{X}{\oplus} \cdots, \qquad (21)$$

where Q^* is the dual of some real vector bundle $Q \to X$ of fiber dimension m. In field models, a vector bundle Q is usually given from the beginning. Therefore, we consider graded manifolds $(X, \mathfrak{A}_Q)$ where the above mentioned isomorphism holds, and call $(X, \mathfrak{A}_Q)$ the simple graded manifold constructed from Q. The structure ring $\mathcal{A}_Q$ of sections of $\mathfrak{A}_Q$ consists of sections of the exterior bundle (21) called graded functions. Given bundle coordinates (x^λ, q^a) on Q with transition functions $q'^a = \rho^a_b q^b$, let $\{c^a\}$ be the corresponding fiber bases for $Q^* \to X$, together with transition functions $c'^a = \rho^a_b c^b$. Then (x^λ, c^a) is called the local basis for the graded manifold $(X, \mathfrak{A}_Q)$. With respect to this basis, graded functions read

$$f = \sum_{k=0}^{m} \frac{1}{k!} f_{a_1\ldots a_k} c^{a_1} \cdots c^{a_k},$$

where $f_{a_1\cdots a_k}$ are local smooth real functions on X.

Given a graded manifold $(X, \mathfrak{A}_Q)$, let $\mathfrak{d}\mathcal{A}_Q$ be the $\mathcal{A}_Q$-module of $\mathbb{Z}_2$-graded derivations of the $\mathbb{Z}_2$-graded ring of $\mathcal{A}_Q$, i.e.,

$$u(ff') = u(f)f' + (-1)^{[u][f]} f u(f'), u \in \mathfrak{d}\mathcal{A}_Q, \qquad f, f' \in \mathcal{A}_Q,$$

where $[.]$ denotes the Grassmann parity. Its elements are called $\mathbb{Z}_2$-graded (or, simply, graded) vector fields on $(X, \mathfrak{A}_Q)$. Due to the canonical splitting $VQ = Q \times Q$, the vertical tangent bundle $VQ \to Q$ of $Q \to X$ can be provided with the fiber bases $\{\partial_a\}$ which is the dual of $\{c^a\}$. Then a graded vector field takes the local form $u = u^\lambda\partial_\lambda + u^a\partial_a$, where u^λ, u^a are local graded functions. It acts on $\mathcal{A}_Q$ by the rule

$$u(f_{a\ldots b}c^a\cdots c^b) = u^\lambda\partial_\lambda(f_{a\ldots b})c^a\cdots c^b + u^d f_{a\ldots b}\partial_d\rfloor(c^a\cdots c^b). \tag{22}$$

This rule implies the corresponding transformation law

$$u'^\lambda = u^\lambda, \qquad u'^a = \rho^a_j u^j + u^\lambda\partial_\lambda(\rho^a_j)c^j.$$

Then one can show [7, 8] that graded vector fields on a simple graded manifold can be represented by sections of the vector bundle $\mathcal{V}_Q \to X$ which is locally isomorphic to the vector bundle $\wedge Q^* \otimes_X (Q \oplus_X TX)$.

Using this fact, we can introduce graded exterior forms on the graded manifold $(X, \mathfrak{A}_Q)$ as sections of the exterior bundle $\wedge\, \mathcal{V}^*_Q$, where $\mathcal{V}^*_Q \to X$ is the $\wedge Q^*$-dual of $\mathcal{V}_Q$. Relative to the dual local bases $\{dx^\lambda\}$ for T^*X and $\{dc^b\}$ for Q^*, graded one-forms read

$$\phi = \phi_\lambda dx^\lambda + \phi_a dc^a, \qquad \phi'_a = \rho^{-1}{}^b{}_a\phi_b, \qquad \phi'_\lambda = \phi_\lambda + \rho^{-1}{}^b{}_a\partial_\lambda(\rho^a_j)\phi_b c^j.$$

The duality morphism is given by the interior product

$$u\rfloor\phi = u^\lambda\phi_\lambda + (-1)^{[\phi_a]}u^a\phi_a.$$

Graded exterior forms constitute the bigraded differential algebra (henceforth BGDA) $\mathcal{C}^*_Q$ with respect to the bigraded exterior product $\wedge$ and the exterior differential d. The standard formulae of a BGDA hold.

Since the jet bundle $J^rQ \to X$ of a vector bundle $Q \to X$ is a vector bundle, let us consider the simple graded manifold $(X, \mathfrak{A}_{J^rQ})$ constructed from $J^rQ \to X$. Its local basis is $\{x^\lambda, c^a_\Lambda\}$, $0 \le |\Lambda| \le r$, together with the transition functions

$$c'^a_{\lambda+\Lambda} = d_\lambda(\rho^a_j c^j_\Lambda), \qquad d_\lambda = \partial_\lambda + \sum_{|\Lambda|<r} c^a_{\lambda+\Lambda}\partial^\Lambda_a, \tag{23}$$

where ∂^Λ_a are the duals of c^a_Λ. Let $\mathcal{C}^*_{J^rQ}$ be the BGDA of graded exterior forms on the graded manifold $(X, \mathfrak{A}_{J^rQ})$. A linear bundle morphism $\pi^r_{r-1} : J^rQ \to J^{r-1}Q$ yields the corresponding monomorphism of BGDAs $\mathcal{C}^*_{J^{r-1}Q} \to \mathcal{C}^*_{J^rQ}$. Hence, there is the direct system of BGDAs

$$\mathcal{C}^*_Q \xrightarrow{\pi^{1*}_0} \mathcal{C}^*_{J^1Q} \cdots \xrightarrow{\pi^r_{r-1}{}^*} \mathcal{C}^*_{J^rQ} \longrightarrow \cdots. \tag{24}$$

Its direct limit $\mathcal{C}^*_\infty Q$ consists of graded exterior forms on graded manifolds $(X, \mathfrak{A}_{J^rQ})$, $r \in \mathbb{N}$, modulo the pull-back identification, and it inherits the BGDA operations intertwined by the monomorphisms $\pi^r_{r-1}{}^*$. It is a $C^\infty(X)$-algebra locally generated by the elements $(1, c^a_\Lambda, dx^\lambda, \theta^a_\Lambda = dc^a_\Lambda - c^a_{\lambda+\Lambda}dx^\lambda)$, $0 \le |\Lambda|$.

In order to regard even and odd dynamic variables on the same footing, let $Y \to X$ be hereafter an affine bundle, and let $\mathcal{P}^*_\infty Y \subset \mathcal{O}^*_\infty Y$ be the $C^\infty(X)$-subalgebra of exterior forms whose coefficients are polynomial in the fiber coordinates y^i_Λ on jet bundles $J^rY \to X$. Let us consider the product

$$\mathcal{S}^*_\infty = \mathcal{C}^*_\infty Q \wedge \mathcal{P}^*_\infty Y \tag{25}$$

of graded algebras $\mathcal{C}^*_\infty Q$ and $\mathcal{P}^*_\infty Y$ over their common graded subalgebra $\mathcal{O}^*X$ of exterior forms on X [5]. It consists of the elements

$$\sum_i \psi_i \otimes \phi_i, \qquad \sum_i \phi_i \otimes \psi_i, \qquad \psi \in \mathcal{C}^*_\infty Q, \qquad \phi \in \mathcal{P}^*_\infty Y,$$

modulo the commutation relations

$$\begin{aligned} &\psi \otimes \phi = (-1)^{|\psi||\phi|}\phi \otimes \psi, \qquad \psi \in \mathcal{C}^*_\infty Q, \qquad \phi \in \mathcal{P}^*_\infty Y, \\ &(\psi \wedge \sigma) \otimes \phi = \psi \otimes (\sigma \wedge \phi), \qquad \sigma \in \mathcal{O}^*X. \end{aligned} \tag{26}$$

They are endowed with the total form degree $|\psi| + |\phi|$ and the total Grassmann parity $[\psi]$. Their multiplication

$$(\psi \otimes \phi) \wedge (\psi' \otimes \phi') := (-1)^{|\psi'||\phi|}(\psi \wedge \psi') \otimes (\phi \wedge \phi'). \tag{27}$$

obeys the relation

$$\varphi \wedge \varphi' = (-1)^{|\varphi||\varphi'| + [\varphi][\varphi']}\varphi' \wedge \varphi, \qquad \varphi, \varphi' \in \mathcal{S}^*_\infty,$$

and makes $\mathcal{S}^*_\infty$ (25) into a bigraded $C^\infty(X)$-algebra. For instance, elements of the ring S^0_∞ are polynomials of c^a_Λ and y^i_Λ with coefficients in $C^\infty(X)$.

The algebra $\mathcal{S}^*_\infty$ is provided with the exterior differential

$$d(\psi \otimes \phi) := (d_\mathcal{C}\psi) \otimes \phi + (-1)^{|\psi|}\psi \otimes (d_\mathcal{P}\phi), \qquad \psi \in \mathcal{C}^*_\infty, \qquad \phi \in \mathcal{P}^*_\infty, \tag{28}$$

where $d_\mathcal{C}$ and $d_\mathcal{P}$ are exterior differentials on the differential algebras $\mathcal{C}^*_\infty Q$ and $\mathcal{P}^*_\infty Y$, respectively. It obeys the relations

$$d(\varphi \wedge \varphi') = d\varphi \wedge \varphi' + (-1)^{|\varphi|}\varphi \wedge d\varphi', \qquad \varphi, \varphi' \in \mathcal{S}^*_\infty,$$

and makes $\mathcal{S}^*_\infty$ into a BGDA, which is locally generated by the elements

$$(1, c^a_\Lambda, y^i_\Lambda, dx^\lambda, \theta^a_\Lambda = dc^a_\Lambda - c^a_{\lambda+\Lambda}dx^\lambda, \theta^i_\Lambda = dy^i_\Lambda - y^i_{\lambda+\Lambda}dx^\lambda), \qquad 0 \leq |\Lambda|.$$

Hereafter, let the collective symbols s^A_Λ and θ^A_Λ stand both for even and odd generating elements c^a_Λ, y^i_Λ, θ^a_Λ, θ^i_Λ of the $C^\infty(X)$-algebra $\mathcal{S}^*_\infty$ which, thus, is locally generated by $(1, s^A_\Lambda, dx^\lambda, \theta^A_\Lambda)$, $|\Lambda| \geq 0$. We agree to call elements of $\mathcal{S}^*_\infty$ the graded exterior forms on X.

Similarly to $\mathcal{O}^*_\infty Y$, the BGDA $\mathcal{S}^*_\infty$ is decomposed into $\mathcal{S}^0_\infty$-modules $\mathcal{S}^{k,r}_\infty$ of k-contact and r-horizontal graded forms together with the corresponding projections h_k and h^r. Accordingly, the exterior differential d (28) on $\mathcal{S}^*_\infty$ is split into the sum $d = d_H + d_V$ of the total and vertical differentials

$$d_H(\phi) = dx^\lambda \wedge d_\lambda(\phi), \qquad d_V(\phi) = \theta^A_\Lambda \wedge \partial^\Lambda_A \phi, \qquad \phi \in \mathcal{S}^*_\infty.$$

One can think of the elements

$$L = \mathcal{L}\omega \in \mathcal{S}^{0,n}_\infty, \qquad \delta(L) = \sum_{|\Lambda|\geq 0} (-1)^{|\Lambda|}\theta^A \wedge d_\Lambda(\partial^\Lambda_A L) \in \mathcal{S}^{0,n}_\infty$$

as being a graded Lagrangian and its Euler–Lagrange operator, respectively.

4 BRST symmetry

A graded derivation $\vartheta \in \mathfrak{d}\mathcal{S}^0_\infty$ of the $\mathbb{R}$-ring $\mathcal{S}^0_\infty$ is said to be contact if the Lie derivative $\mathbf{L}_\vartheta$ preserves the ideal of contact graded forms of the BGDA $\mathcal{S}^*_\infty$. With respect to the local basis $(x^\lambda, s^A_\Lambda, dx^\lambda, \theta^A_\Lambda)$ for the BGDA $\mathcal{S}^*_\infty$, any contact graded derivation takes the form

$$\vartheta = \vartheta_H + \vartheta_V = \vartheta^\lambda d_\lambda + (\vartheta^A \partial_A + \sum_{|\Lambda|>0} d_\Lambda \vartheta^A \partial^\Lambda_A), \tag{29}$$

where ϑ^λ, ϑ^A are local graded functions [5]. The interior product $\vartheta\rfloor\phi$ and the Lie derivative $\mathbf{L}_\vartheta\phi$, $\phi \in \mathcal{S}^*_\infty$, are defined by the same formulae

$$\begin{aligned}
&\vartheta\rfloor\phi = \vartheta^\lambda \phi_\lambda + (-1)^{[\phi_A]}\vartheta^A \phi_A, \qquad \phi \in \mathcal{S}^1_\infty, \\
&\vartheta\rfloor(\phi\wedge\sigma) = (\vartheta\rfloor\phi)\wedge\sigma + (-1)^{|\phi|+[\phi][\vartheta]}\phi\wedge(\vartheta\rfloor\sigma), \qquad \phi,\sigma \in \mathcal{S}^*_\infty \\
&\mathbf{L}_\vartheta\phi = \vartheta\rfloor d\phi + d(\vartheta\rfloor\phi), \qquad \mathbf{L}_\vartheta(\phi\wedge\sigma) = \mathbf{L}_\vartheta(\phi)\wedge\sigma + (-1)^{[\vartheta][\phi]}\phi\wedge\mathbf{L}_\vartheta(\sigma).
\end{aligned}$$

as those on a graded manifold. One can justify that any vertical contact graded derivation ϑ (29) satisfies the relations

$$\vartheta\rfloor d_H\phi = -d_H(\vartheta\rfloor\phi), \qquad \mathbf{L}_\vartheta(d_H\phi) = d_H(\mathbf{L}_\vartheta\phi), \qquad \phi \in \mathcal{S}^*_\infty. \tag{30}$$

Proposition 2. The Lie derivative $\mathbf{L}_\vartheta L$ of a Lagrangian L along a contact graded derivation ϑ (29) fulfills the first variational formula

$$\mathbf{L}_\vartheta L = \vartheta_V\rfloor\delta L + d_H(h_0(\vartheta\rfloor\Xi_L)) + d_V(\vartheta_H\rfloor\omega)\mathcal{L}, \tag{31}$$

where $\Xi_L = \Xi + L$ is a Lepagean equivalent of a graded Lagrangian L [5].

A contact graded derivation ϑ is said to be variational if the Lie derivative (31) is d_H-exact. A glance at the expression (31) shows that: (i) A contact graded derivation ϑ is variational only if it is projected onto X, and (ii) ϑ is variational iff its vertical part ϑ_V is well. Therefore, we restrict our consideration to vertical contact graded derivations

$$\vartheta = \sum_{0\leq|\Lambda|} d_\Lambda \upsilon^A \partial_A^\Lambda. \tag{32}$$

Such a derivation is completely defined by its first summand

$$\upsilon = \upsilon^A(x^\lambda, s^A_\Lambda)\partial_A, \qquad 0 \leq |\Lambda| \leq k, \tag{33}$$

which is also a graded derivation of $\mathcal{S}^0_\infty$. It is called the generalized graded vector field. A glance at the first variational formula (31) shows that ϑ (32) is variational iff $\upsilon\rfloor\delta L$ is d_H-exact.

A vertical contact graded derivation ϑ (32) is said to be nilpotent if

$$\mathbf{L}_\upsilon(\mathbf{L}_\upsilon\phi) = \sum_{|\Sigma|\geq 0,|\Lambda|\geq 0} (\upsilon^B_\Sigma \partial^\Sigma_B(\upsilon^A_\Lambda)\partial^\Lambda_A + (-1)^{[s^B][\upsilon^A]}\upsilon^B_\Sigma \upsilon^A_\Lambda \partial^\Sigma_B \partial^\Lambda_A)\phi = 0 \tag{34}$$

for any horizontal graded form $\phi \in S^{0,*}_\infty$ or, equivalently, $\vartheta\circ\vartheta)(f) = 0$ for any graded function $f \in \mathcal{S}^0_\infty$. One can show that ϑ is nilpotent only if it is odd and iff the equality

$$\vartheta(\upsilon^A) = \sum_{|\Sigma|\geq 0} \upsilon^B_\Sigma \partial^\Sigma_B(\upsilon^A) = 0 \tag{35}$$

holds for all υ^A [5].

Return now to the original gauge system on a fiber bundle Y with a Lagrangian L (10) and a gauge symmetry u (3). For the sake of simplicity, $Y \to X$ is assumed to be affine. Let us consider the BGDA $\mathcal{S}^*_\infty = \mathcal{C}^*_\infty V \wedge \mathcal{P}^*_\infty Y$ locally generated by $(1, c^r_\Lambda, dx^\lambda, y^i_\Lambda, \theta^r_\Lambda, \theta^i_\Lambda)$. Let $L \in \mathcal{O}^{0,n}_\infty Y$ be a polynomial in y^i_Λ, $0 \leq |L|$. Then it is a graded Lagrangian $L \in \mathcal{P}^{0,n}_\infty Y \subset \mathcal{S}^{0,n}_\infty$ in $\mathcal{S}^*_\infty$. Its gauge symmetry u (3) gives rise to the generalized vector field $u_E = u$ on E, and the latter defines the generalized graded vector field υ (33) by the formula (5). It is easily justified that the contact graded derivation ϑ (32) generated by υ (5) is variational for L. It is odd, but need not be nilpotent. However, one can try to find a nilpotent contact graded derivation (32) generated by some generalized graded vector field (6) which coincides with ϑ on $\mathcal{P}^*_\infty Y$. We agree to call it the BRST operator.

In this case, the nilpotency conditions (35) read

$$\sum_\Sigma d_\Sigma(\sum_\Xi \upsilon^{i,\Xi}_r c^r_\Xi) \sum_\Lambda \partial^\Sigma_i(\upsilon^{j,\Lambda}_s)c^s_\Lambda + \sum_\Lambda d_\Lambda(u^r)\upsilon^{j,\Lambda}_r = 0, \tag{36}$$

$$\sum_\Lambda(\sum_\Xi d_\Lambda(\upsilon^{i,\Xi}_r c^r_\Xi)\partial^\Lambda_i + d_\Lambda(u^r)\partial^\Lambda_r)u^q = 0 \tag{37}$$

for all indices j and q. They are equations for graded functions $u^r \in \mathcal{S}^0_\infty$. Since these functions are polynomials

$$u^r = u^r_{(0)} + \sum_\Gamma u^{r,\Gamma}_{(1)p} c^p_\Gamma + \sum_{\Gamma_1,\Gamma_2} u^{r,\Gamma_1\Gamma_2}_{(2)p_1p_2} c^{p_1}_{\Gamma_1} c^{p_2}_{\Gamma_2} + \cdots \tag{38}$$

in c^s_Λ, the equations (36) – (37) take the form

$$\sum_\Sigma d_\Sigma(\sum_\Xi \upsilon^{i,\Xi}_r c^r_\Xi) \sum_\Lambda \partial^\Sigma_i(\upsilon^{j,\Lambda}_s) c^s_\Lambda + \sum_\Lambda d_\Lambda(u^r_{(2)}) \upsilon^{j,\Lambda}_r = 0, \tag{39}$$

$$\sum_\Lambda d_\Lambda(u^r_{(k\neq 2)}) \upsilon^{j,\Lambda}_r = 0, \tag{40}$$

$$\sum_\Lambda \sum_\Xi d_\Lambda(\upsilon^{i,\Xi}_r c^r_\Xi) \partial^\Lambda_i u^q_{(k-1)} + \sum_{m+n-1=k} d_\Lambda(u^r_{(m)}) \partial^\Lambda_r u^q_{(n)} = 0. \tag{41}$$

If the equations (39) – (41) have a solution, i.e., the (nilpotent) BRST operator exists, one can think of the equalities (39) and (41) (and, consequently, the nilpotency conditions (36) – (37)) as being the generalized commutation relations and generalized Jacobi identities of gauge transformations, respectively.

Indeed, the relation (39) for components υ^i_r takes the form of the familiar Lie bracket

$$\sum_\Sigma [d_\Sigma(\upsilon^i_p) \partial^\Sigma_i \upsilon^j_q - d_\Sigma(\upsilon^i_q) \partial^\Sigma_i \upsilon^j_p] = -2u^r_{(2)pq} \upsilon^j_r,$$

where $-2u^r_{(2)pq}$ are generalized structure constants depending on dynamic variables y^i and their jets y^i_Λ in general. For instance, let us assume that all υ^i_r are linear in y^i_Λ. Then $u^r_{(2)pq}$ are independent of these variables. Let $u^r_{(m\neq 2)} = 0$. In this case, the relation (41) reduces to the familiar Jacobi identity

$$u^r_{(2)pq} u^j_{(2)rs} + u^r_{(2)qs} u^j_{(2)rp} + u^r_{(2)sp} u^j_{(2)rq} = 0.$$

Let us note that any Lagrangian L have gauge symmetries. In particular, there always exist trivial gauge symmetries

$$\upsilon = \sum_\Lambda T^{j,i,\Lambda}_r \mathcal{E}_j \xi^r_\Lambda \partial_i, \qquad T^{j,i,\Lambda}_r = -T^{i,j,\Lambda}_r,$$

vanishing on-shell. In a general setting, one therefore can require that the nilpotency conditions (36) – (37) hold on-shell, i.e., gauge transformations form an algebra on-shell.

5 Example

Let us consider the gauge theory of principal connections on a principal bundle $P \to X$ with a structure Lie group G. These connections are represented by sections of the quotient

$$C = J^1P/G \to X. \tag{42}$$

This is an affine bundle coordinated by (x^λ, a^r_λ) such that, given a section A of $C \to X$, its components $A^r_\lambda = a^r_\lambda \circ A$ are coefficients of the familiar local connection form (i.e., gauge potentials). Let $J^\infty C$ be the infinite order jet manifold of $C \to X$ coordinated by $(x^\lambda, a^r_{\Lambda\lambda})$, $0 \leq |\Lambda|$. We consider the GDA $\mathcal{O}^*_\infty C$.

Infinitesimal generators of one-parameter groups of automorphisms of a principal bundle P are G-invariant projectable vector fields on $P \to X$. They are associated to sections of the vector bundle $T_GP = TP/G \to X$. This bundle is provided with the coordinates $(x^\lambda, \dot{x}^\lambda, \xi^r)$ with respect to the fibre bases $\{\partial_\lambda, e_r\}$ for T_GP, where $\{e_r\}$ is the basis for the right Lie algebra $\mathfrak{g}$ of G such that $[e_p, e_q] = c^r_{pq} e_r$. If

$$u = u^\lambda \partial_\lambda + u^r e_r, \qquad v = v^\lambda \partial_\lambda + v^r e_r, \tag{43}$$

are sections of $T_GP \to X$, their bracket reads

$$[u, v] = (u^\mu \partial_\mu v^\lambda - v^\mu \partial_\mu u^\lambda)\partial_\lambda + (u^\lambda \partial_\lambda v^r - v^\lambda \partial_\lambda u^r + c^r_{pq} u^p v^q) e_r. \tag{44}$$

Any section u of the vector bundle $T_GP \to X$ yields the vector field

$$u_C = u^\lambda \partial_\lambda + (c^r_{pq} a^p_\lambda u^q + \partial_\lambda u^r - a^r_\mu \partial_\lambda u^\mu)\partial^\lambda_r \tag{45}$$

on the bundle of principal connections C (42). It is an infinitesimal generator of a one-parameter group of automorphisms of C [7]. Let us consider the bundle product

$$E = C \underset{X}{\times} T_GP, \tag{46}$$

coordinated by $(x^\lambda, \tau^\lambda = \dot{x}^\lambda, \xi^r, a^r_\lambda)$. It can be provided with the generalized vector field

$$\upsilon_E = \upsilon = (c^r_{pq} a^p_\lambda \xi^q + \xi^r_\lambda - a^r_\mu \tau^\mu_\lambda - \tau^\mu a^r_{\mu\lambda})\partial^\lambda_r. \tag{47}$$

Following the procedure in Sections 3 – 4, we replace parameters ξ^r and τ^λ with the odd ghosts c^r and c^λ, respectively, and obtain the generalized graded vector field

$$\upsilon = (c^r_{pq} a^p_\lambda c^q + c^r_\lambda - a^r_\mu c^\mu_\lambda - c^\mu a^r_{\mu\lambda})\partial^\lambda_r + (-\frac{1}{2} c^r_{pq} c^p c^q - c^\mu c^r_\mu)\partial_r + c^\lambda_\mu c^\mu \partial_\lambda \tag{48}$$

such that the vertical contact graded derivations (32) generated by υ (48) is nilpotent, i.e., it is a BRST operator.

References

[1] G.Barnich, F.Brandt and M.Henneaux, Local BRST cohomology in gauge theories, *Phys. Rep.* **338**, 439 (2000).

[2] D.Bashkirov, G.Giachetta, L.Mangiarotti and G.Sardanashvily, Noether's second theorem in a general setting. Reducible gauge theories, *E-print arXiv*: math.DG/0411070.

[3] F.Brandt, Jet coordinates for local BRST cohomology, *Lett. Math. Phys.* **55**, 149 (2001).

[4] R.Fulp, T.Lada and J.Stasheff, Sh-Lie algebras induced by gauge transformations, *Comm. Math. Phys.* **231**, 25 (2002); *E-print arXiv*: math.QA/0012106.

[5] G.Giachetta, L.Mangiarotti and G.Sardanashvily, Lagrangian supersymmetries depending on derivatives. Global analysis and cohomology, *Commun. Math. Phys.* (accepted); *E-print arXiv*: hep-th/0407185.

[6] J.Gomis, J.París, J. and S.Samuel, Antibracket, antifields and gauge theory quantization, *Phys. Rep* **295**, 1 (1995).

[7] L.Mangiarotti and G.Sardanashvily, *Connections in Classical and Quantum Field Theory* (World Scientific, Singapore, 2000).

[8] G.Sardanashvily, SUSY-extended field theory, *Int. J. Mod. Phys. A* **15**, 3095 (2000).